

EHHI BOARD MEMBERS

Susan S. Addiss, MPH, MUR.S. Past Commissioner of Health for the State of Connecticut; Past President of the American Public Health Association; Past Member of the Pew Environmental Health Commission; Vice-Chair, Connecticut Health Foundation Board; Director of Health Education for Environment and Human Health, Inc.

Nancy O. Alderman, MES. President of Environment and Human Health, Inc.; Past Member of the Governor's Pollution Prevention Task Force; Past Member of the National Board of Environmental Defense; Recipient of the Connecticut Bar Association, Environmental Law Section's, Clyde Fisher Award, given in recognition of significant contributions to the preservation of environmental quality through work in the fields of environmental law, environmental protection or environmental planning, and the New England Public Health Association's Robert C. Huestis/Eric Mood Award given to individuals for outstanding contributions to public health in the environmental health area.

D. Barry Boyd, M.D. Clinical Professor of Medicine, Yale University School of Medicine; Oncologist at Greenwich Hospital and Affiliate Member of the Yale Cancer Center. Research areas include environmental risk factors for cancer as well as cancer etiology, including nutrition and the role of insulin and IGF in malignancy. Dr. Boyd is the Founder and Director of Integrative Medicine at Greenwich Hospital – Yale Health System.

Russell L. Brenneman, Esq. Environmental lawyer and activist; Founding Co-Chair of the Connecticut League of Conservation Voters; Past Chair of the CTLCV Education Fund; Adjunct faculty, Trinity College, Hartford, Connecticut; Past President, Connecticut Forest and Park Association.

David R. Brown, Sc.D. Public Health Toxicologist and Director of Public Health Toxicology for Environment and Human Health, Inc.; Past Chief of Environmental Epidemiology and Occupational Health at Connecticut's Department of Health; Past Deputy Director of The Public Health Practice Group of ATSDR at the national Centers for Disease Control and Prevention (CDC) in Atlanta, Georgia.

Robert G. LaCamera, M.D. Clinical Professor of Pediatrics, Yale University School of Medicine; Primary Care Pediatrician in New Haven, Connecticut from 1956 to 1996 with a sub-specialty in children with disabilities.

Peter M. Rabinowitz, M.D., Ph.D. Associate Professor of Occupational and Environmental Medicine, Yale University School of Medicine. Director of clinical services at Yale's Department of Occupational and Environmental Medicine. Principal investigator on the Canary Database Project, which looks at animals as sentinels of environmental health hazards.

Hugh S. Taylor, M.D. Professor of Obstetrics, Gynecology and Reproductive Sciences and Department of Molecular, Cellular and Developmental Biology; Chief of the Division of Reproductive Endocrinology and Infertility, Yale University School of Medicine.

John P. Wargo, Ph.D. Professor of Risk Analysis and Environmental Policy, Yale University's School of Forestry and Environmental Studies, and Professor of Political Science. Director of Yale College's Environmental Studies major. Director of the Yale Program on Environment and Health; Author of *Our Children's Toxic Legacy* and *Green Intelligence*.

12 Steps to Reducing Fetal Exposures



Although obstetricians warn pregnant women not to drink alcohol and not to smoke during pregnancy, we are learning that there are other chemicals and products that women should be careful about during those critical nine months. The Environmental Working Group found more than 200 chemicals in the cord blood of babies tested right after birth. Below are twelve suggestions of ways that may help reduce some of these exposures.

1. Pesticides

The safest policy is to avoid using pesticides in your home or garden during pregnancy. Avoid them particularly during the first trimester when the baby's neural tube and nervous system are developing. The California Birth Defects Monitoring Program reported that three out of every four women are exposed to pesticides around the home. They also observed that



pregnant women exposed to household gardening pesticides had a modestly increased risk of having babies with oral clefts, neural tube defects, heart defects, and limb defects. The journal

Environmental Health Perspectives (EHP), Volume 110, reports that children who are exposed to indoor pesticides are at an elevated risk for leukemia. EHP adds that the risk is increased during the first three months of pregnancy and when professional pest control services are used in the home.

2. Paint

The current recommendation is to avoid exposure to oil-based paints, lead and mercury. However, you



should also limit exposure to latex paints that contain ethylene glycol ethers and biocides. Biocides are used to retard mold. Ideally, you should get someone else to do the job for you. Recent studies show

that infants and children who sleep in bedrooms with fumes from water-based paints and solvents are two to four times more likely to suffer allergies and asthma. It would be wise, therefore, to reduce exposures during pregnancy as well.



ENVIRONMENT & HUMAN HEALTH, INC.

1191 Ridge Road

North Haven, Connecticut 06473

Phone (203) 248-6582 Fax (203) 288-7571

www.ehhi.org

3. Solvents, paint thinners, paint strippers, polyurethane, etc.

The *Journal of the American Medical Association* reports significantly more malformations in the fetuses of mothers who were occupationally exposed to solvents. Turpentine, liquid paint removers, and other solvents should all be avoided during pregnancy. Polyurethaning hardwood floors should also be avoided.



4. Toxic cleaning products

Oven cleaners and strong tile cleaners — both of which are used in tight spaces — should be avoided. Any cleaning product with a strong chemical odor should be avoided. If you do use chemical cleaning products, make sure that you have good ventilation and that you wear gloves and clothing to protect your skin. Always read the warning and instruction labels. Never mix different chemicals together because the combination may create poisonous fumes, as when you mix ammonia and bleach.



5. Fragrances

A study of selected scented consumer products showed that they emitted more than 100 volatile organic compounds (VOCs). Some of them are classified as toxic or hazardous chemicals according to federal laws. Ten of the 100 VOCs identified qualify under federal rules as toxic or hazardous. Some air fresheners plug into electrical sockets and pulsate air freshener pollutants on a continual basis. Scented candles also emit VOCs and are scented with synthetic oils that provide additional sources of petroleum-based fumes. Those most at risk from the fumes released by scented candles are pregnant women, children, and pets.



6. Plastics: Phthalates and Bisphenol A

Bisphenol A (BPA) is estrogenic and has been shown to cause heart disease, obesity, diabetes, and reproductive health problems in experiments with laboratory animals. BPA is used in a wide variety of consumer products and can be found in the linings of tin cans, and in hard and clear plastic containers that often have the number 7 or the letters “PC” for the chemical polycarbonate on the bottom. Phthalates, which are endocrine disruptors, are used to soften plastics such as plastic shower curtains, some children’s



toys and vinyl flooring. Phthalates are also used to enhance fragrances and are frequently found in scented products. They disrupt the endocrine system by mimicking testosterone. Fetal exposures have been shown to cause a variety of reproductive abnormalities in rats.

7. Teflon-coated cooking utensils

A study done in 2005 by the nonprofit organizations Environmental Working Group and Commonwealth found perfluorooctanoic acid (PFOA) in the umbilical cord blood of newborns. Johns Hopkins Medical Center, which did a similar test in 2006, found PFOA present in the umbilical cord blood of 99% of the 300 infants tested. The U.S. EPA has declared PFOA a possible carcinogen. PFOA is a chemical found in teflon-coated pans and stain-resistant fabrics.



8. Cell phones



Pregnant women should **not** keep cell phones that are turned on close to their bodies. It is recommended when using a cell phone to either put it on speaker mode or use a wired earpiece.

9. Cat litter

Cats are carriers of the parasite toxoplasmosis, which lives inside their intestines and is passed on in their feces. Therefore, it is recommended that pregnant women wear gloves, a mask, and that they wash their hands thoroughly when cleaning cat litter. The effects of toxoplasmosis are mild to the mother, but not to the fetus. Toxoplasmosis can cause both vision and hearing loss, as well as developmental delays in the affected child.



10. Fish containing mercury

The American Pregnancy Association recommends that pregnant women (1) **avoid** eating swordfish, tuna (bigeye, ahi); and (2) **eat no more than three 6-oz servings per month** of sea bass (Chilean), bluefish, grouper, tuna (canned, white albacore), tuna (yellowfin); and (3) **eat no more than six 6-oz servings per month** of bass (striped, black), cod (Alaskan), halibut (Pacific and Atlantic), lobster, mahi-mahi, and snapper.



11. Artificial sweeteners

During pregnancy, women are advised to use artificial sweeteners in moderation. However, studies show that saccharin (Sweet 'N Low) crosses the placenta and may remain in fetal tissue. Therefore, its use by pregnant women remains in question.



12. Clean air and adequate ventilation

Clean and well-ventilated air is important for everyone’s good health. Air pollutants should be kept at a minimum, both inside and outside the home and workplace. This is especially important during pregnancy.

